

Quantum Corrections to Entropy of Charged Dilatonic Black Holes in Arbitrary Dimensions

Kiyoshi Shiraishi

Akita Junior College, Shimokitade-Sakura, Akita-shi,
Akita 010, Japan

Modern Physics Letters **A9** (1994) pp. 3509-3516

Abstract

The quantum contribution of a scalar field to entropy of a dilatonic black hole is calculated in the brick wall model by the WKB method and analyzed by a high-temperature expansion. If the cutoff distance from the horizon approaches zero, the leading divergent piece of entropy turns out to be proportional to the “area” of the horizon surface (which has $(N - 1)$ -dimensional extension in $(N + 1)$ -dimensional space-time) and independent of other properties of black holes even in the case of general dilaton coupling. There is also qualitative argument with the known result of subleading divergence for $N = 3$.

PACS Nos.: 04.60.+n, 12.10.Gq, 97.60.Lf, 11.10.Kk.

Recently, there has been a renewed interest in thermodynamic description of black holes. The “zero-loop” partition function of black holes is derived from the path-integral of a classical action of Einstein gravitation, along with a special attention to the boundary conditions.[1]

An attempt to calculate contribution of quantum fields to the black hole entropy was made in the seminal work of 't Hooft.[2] In the last decade, entropy due to quantum fields and string excitations has been investigated by many authors. Recently, Callan and Wilzcek analyzed field theory in an accelerated system and obtained quantum contribution to the entropy, which they called as geometric entropy.[3] Their geometric entropy is believed to be equivalent to the one obtained in the other ways such as was done in Ref. [4]. Susskind and Uglum [5] and Kabat and Strasslers [6] showed that similar result can be obtained through the WKB method, which is used in the 't Hooft's analysis.[2]

In a recent paper,[7] Ghosh and Mitra calculated the entropy due to quantum scalar field in the background of a dilatonic black hole in effective string theory by the WKB approximation. They showed the cutoff-dependent part is not proportional to the area of the horizon surface if the black hole is maximally charged. Recent calculation by Solodukhin [8] supports their evaluation.

In this letter, we compute quantum corrections to black hole entropy in a model with an arbitrary dilaton coupling in arbitrary space-time dimensions using the WKB method. First of all, we start with reviewing the classical background of a charged dilatonic black hole. Suppose the classical action for gravity with a U(1) gauge field A_μ and a dilaton field as follows:

$$S = \int d^{N+1}x \frac{\sqrt{-g}}{16\pi} \left[R - \frac{4}{N-1} (\nabla\phi)^2 - e^{-4a\phi/(N-1)} F^2 \right] + (\text{boundary terms}), \quad (1)$$

where $F^2 = F^{\mu\nu} F_{\mu\nu}$ with the field strength $F_{\mu\nu} = \partial_\mu A_\nu - \partial_\nu A_\mu$ and the Newton constant is set to unity. Here we assume that the dilaton coupling a may take any positive value: For effective field theories of string theory, it takes $a = 1$.

The spherical black hole solution to the field equation derived from the action (1) is given by [9, 10]

$$ds^2 = -\sigma^{-2}(r) \Delta(r) dt^2 + \sigma^{2/(N-2)}(r) \left[\frac{dr^2}{\Delta(r)} + r^2 d\Omega_{N-1}^2 \right], \quad (2)$$

with

$$\Delta(r) = \left[1 - \left(\frac{r_+}{r} \right)^{N-2} \right] \left[1 - \left(\frac{r_-}{r} \right)^{N-2} \right], \quad (3)$$

$$\Delta(r) = \left[1 - \left(\frac{r_-}{r} \right)^{N-2} \right]^{a^2/(N-2+a^2)}, \quad (4)$$

where r_+ and r_- are constants ($r_+ > r_-$). In Eq. (2), $d\Omega_{N-1}^2$ represents the line element of a unit $(N-1)$ -sphere.

For this solution, the dilaton field configuration is expressed as

$$e^{4a\phi/(N-1)} = \sigma^2(r) \quad (5)$$

and the electric field is written as

$$F = \frac{Q}{r^{N-1}} dt \wedge dr, \quad (6)$$

where the charge Q is related to r_+ and r_- as

$$Q^2 = \frac{(N-1)(N-2)^2}{2(N-2+a^2)} (r_+ r_-)^{N-2}. \quad (7)$$

The mass M of the black hole described by the solution is given by

$$M = \frac{(N-1)A_{N-1}}{16\pi} \left(r_+^{N-2} + \frac{N-2-a^2}{N-2+a^2} r_-^{N-2} \right), \quad (8)$$

where

$$A_{N-1} = \frac{2\pi^{N/2}}{\Gamma(N/2)}. \quad (9)$$

Now we introduce an extra neutral scalar field φ with a mass m in the black hole background. The scalar is allowed to be coupled with the background dilaton field in general. We will evaluate the contribution of this field to black hole entropy by means of a mode function technique. The equation of motion of the scalar field can be written in the general form:

$$\nabla^\mu e^{-4ab\phi/(N-1)} \nabla_\mu \varphi = e^{-4ac\phi/(N-1)} m^2 \varphi, \quad (10)$$

where b and c are parameters representing the strength of the coupling to the dilaton field.

In the spherically symmetric space, the scalar field can be written by a product of mode functions:

$$\varphi = \frac{1}{r^{(N-1)/2}} e^{-i\omega t} \chi_{\omega\ell}(r) Y_{\ell\nu}^{(N-1)}(\Omega), \quad (11)$$

where the generalized spherical function $Y_{\ell\nu}^{(N-1)}(\Omega)$ is the eigenfunction of the Laplacian on S^{N-1} with a unit radius, such that

$$\Delta^{(N-1)} Y_{\ell\nu}^{(N-1)} = -\ell(\ell + N - 2) Y_{\ell\nu}^{(N-1)}, \quad (\ell = 0, 1, 2, \dots). \quad (12)$$

The differential equation concerning the radial function $\chi_{\omega\ell}(r)$ can be obtained by substituting Eq. (11) into Eq. (10). We further introduce a new radial coordinate y defined by [10]

$$y = \int^r \frac{\sigma^{2b}}{\Delta} dr, \quad (13)$$

where y varies between $-\infty$ and ∞ while r varies between r_+ and ∞ . Then the wave equation for the radial function is simplified as [10]

$$\frac{d^2 \chi_{\omega\ell}}{dy^2} + W \chi_{\omega\ell} = 0, \quad (14)$$

where

$$W(y) = \left(\frac{\Delta(r)}{\sigma^{2b}(r)} \right)^2 \frac{\sigma^{2/(N-2)}(r)}{\Delta(r)} \left[\frac{\sigma^2(r)}{\Delta(r)} \omega^2 - \frac{\ell(\ell + N - 2)}{r^2 \sigma^{2/(N-2)}(r)} \right. \\ \left. - \frac{N-1}{2r \sigma^{2/(N-2)}(r)} \frac{d\Delta(r)}{dr} - m^2 \sigma^{2(b-c)}(r) - \Delta(r) \mu^2(r) \right], \quad (15)$$

with

$$\mu^2(r) = \frac{1}{\sigma^{2/(N-2)}(r)} \left[\frac{(N-1)(N-3)}{4r^2} - \frac{N-1}{r} b \frac{1}{\sigma(r)} \frac{d\sigma(r)}{dr} \right]. \quad (16)$$

Let us use the WKB method to evaluate the solution to Eq. (14). The solution is approximated as

$$\chi_{\omega\ell}(y) \approx \frac{1}{\sqrt{W(y)}} \int^y \sin(\sqrt{W(y)}) dy. \quad (17)$$

If we impose the boundary condition that the wave function vanishes on the “brick wall” at $r = r_+ + \delta$ and on the wall of a large box at $r = L$, [2] we find that the discrete quantum number n is associated with the phase function [2, 5, 6, 7, 11]:

$$n\pi = \int_{r_+ + \delta}^L k_{\omega\ell}(r) dr, \quad (18)$$

where

$$k_{\omega\ell}(r) = \sqrt{\frac{\sigma^{2/(N-2)}(r)}{\Delta(r)} \left[\frac{\sigma^2}{\Delta} \omega^2 - \frac{\ell(\ell + N - 2)}{r^2 \sigma^{2/(N-2)}} - \frac{N - 1}{2r \sigma^{2/(N-2)}} \frac{d\Delta}{dr} - m^2 \sigma^{2(b-c)} - \Delta\mu^2 \right]}. \quad (19)$$

The free energy F can be derived from the density of the modes. After some manipulation, [2, 5, 6, 7, 11] one can get the following expression for the free energy at inverse temperature β :

$$F = -\frac{1}{\pi} \sum_{\ell=0}^{\infty} D_{\ell} \int d\omega \frac{1}{e^{\beta\omega} - 1} \int_{r_+ + \delta}^L k_{\omega\ell}(r) dr \equiv \int_{r_+ + \delta}^L f(r) dr, \quad (20)$$

where the degeneracy of the spherical modes D_{ℓ} is [12]

$$D_{\ell} \equiv \frac{(2\ell + N - 2)(\ell + N - 3)!}{(N - 2)! \ell!}. \quad (21)$$

In (20), the integral over ω should be taken only over possible values for which $k_{\omega\ell}$ is real.

We wish to study the portion that yields the divergence of the free energy as $\delta \rightarrow 0$. In the first step to this end, we approximate the “density” $f(r)$. Integrating over ω , $f(r)$ can be rewritten by using the modified Bessel function [13] as

$$f(r) = -\frac{\sigma^{-(N-3)/(N-2)}}{\pi} \sum_{\ell=0}^{\infty} D_{\ell} \sum_{p=1}^{\infty} \frac{\eta(r)}{p\beta\sqrt{\Delta\sigma^{-2}}} K_1(p\beta\sqrt{\Delta\sigma^{-2}}\eta(r)), \quad (22)$$

where $\eta(r)$ is defined by

$$\eta(r) = \sqrt{\frac{\ell(\ell + N - 2)}{r^2 \sigma^{2/(N-2)}} + \frac{N - 1}{2r \sigma^{2/(N-2)}} \frac{d\Delta}{dr} + m^2 \sigma^{2(b-c)} + \Delta\mu^2}. \quad (23)$$

Note that the inverse temperature β appears only in the combination $\beta\sqrt{\Delta\sigma^{-2}} = \beta\sqrt{|g_{tt}|} \equiv \tilde{\beta}(r)$. To see the structure of generating divergences in free energy, we will expand Eq. (22) with respect to powers of $\tilde{\beta}(r)$; this is just a “high-temperature” expansion. To this end, we use another integral representation of the modified Bessel function. [13] Then Eq. (22) becomes

$$f(r) = -\frac{\sigma^{-(N-3)/(N-2)}}{\pi} \sum_{\ell=0}^{\infty} D_{\ell} \sum_{p=1}^{\infty} \frac{1}{p^2 \tilde{\beta}^2} \int_0^{\infty} \exp\left(-t - \frac{p^2 \tilde{\beta}^2 \eta^2}{4t}\right) dt. \quad (24)$$

To perform the sum over ℓ , we use the following expansion [12] in Eq. (24):

$$\sum_{\ell=0}^{\infty} D_{\ell} e^{-\ell(\ell+N-2)x} = \frac{\Gamma((N-1)/2)}{\Gamma(N-1)} x^{-(N-1)/2} \left[1 + \frac{(N-1)(N-2)}{6} x + O(x^2) \right]. \quad (25)$$

Consequently, we find the following expression to the next-to-leading order for small $\tilde{\beta}(r)$, under the assumption of $m\sigma^{b-c} \ll \tilde{\beta}^{-1}$ and $r\sigma^{1/(N-1)} \gg \tilde{\beta}$:

$$\begin{aligned} f(r) = & -\frac{A_{N-1} r^{N-1} \sigma^{2/(N-2)}}{\pi^{(N+1)/2}} \left\{ \frac{1}{\tilde{\beta}^{N+1}} \Gamma((N+1)/2) \zeta(N+1) \right. \\ & + \frac{1}{4\tilde{\beta}^{N-1}} \Gamma((N-1)/2) \zeta(N-1) \left[\frac{(N-1)(N-2)}{6r^2 \sigma^{2/(N-2)}} - \frac{N-1}{2r \sigma^{2/(N-2)}} \frac{d\Delta}{dr} \right. \\ & \left. \left. - m^2 \sigma^{2(b-c)} - \Delta \mu^2 \right] \right\} + O(\tilde{\beta}^{-(N-3)}), \end{aligned} \quad (26)$$

where $\zeta(z)$ is the Riemann zeta function and we have applied formulas for the gamma functions [13].

In the second step, we examine the divergence of free energy after integration over r as $\delta \rightarrow 0$. We first focus our attention on the contribution from the leading term of the high-temperature expansion.

The estimation can be done in a straightforward way. To rearrange the result, we prepare the following quantities. The invariant distance between the horizon at $r = r_+$ and $r = r_+ + \delta$ is [2]

$$\begin{aligned} \varepsilon = & 2 \left(1 - \frac{r_-^{N-2}}{r_+^{N-2}} \right)^{a^2/[(N-2)(N-2+a^2)]-1/2} \sqrt{\frac{r_+ \delta}{N-2}} \\ & \times \left\{ 1 + \frac{1}{6} \frac{\delta}{r_+} \left[\frac{N-1}{2} + \left(\frac{2a^2}{N-2+a^2} - (N-2) \right) \frac{\frac{r_-^{N-2}}{r_+^{N-2}}}{1 - \frac{r_-^{N-2}}{r_+^{N-2}}} \right] + O((\delta/r_+)^2) \right\} \end{aligned} \quad (27)$$

The Hawking temperature of the dilatonic black hole is given by [9, 10]

$$T_H = \beta_H^{-1} = \frac{N-2}{4\pi r_+} \left(1 - \frac{r_-^{N-2}}{r_+^{N-2}} \right)^{[(N-2)^2 - a^2]/[(N-2)(N-2+a^2)]}. \quad (28)$$

Then we find that the leading term of the divergent part as $\delta \rightarrow 0$, or equivalently as $\varepsilon \rightarrow 0$, in the free energy of the order of $\beta^{-(N+1)}$ is expressed as:

$$F_{div1} = -\frac{1}{(N-1)(2\pi)^N} \frac{\Gamma((N+1)/2) \zeta(N+1)}{\pi^{(N+1)/2}} \frac{\tilde{A}_H^{(N-1)}}{\varepsilon^{N-1}} \frac{\beta_H^N}{\beta^{N+1}}, \quad (29)$$

where the “area” of the horizon $(N-1)$ -sphere is given by

$$\tilde{A}_H^{(N-1)} \equiv A_{N-1} \tilde{r}_+^{N-1}, \quad (30)$$

with

$$\tilde{r}_+ \equiv r_+ \sigma^{1/(N-2)}(r_+). \quad (31)$$

The corresponding entropy turns out to be

$$S_{div1} = \beta^2 \frac{\partial F_{div1}}{\partial \beta} = \frac{N+1}{(N-1)(2\pi)^N} \frac{\Gamma((N+1)/2)\zeta(N+1)}{\pi^{(N+1)/2}} \frac{\tilde{A}_H^{(N-1)}}{\varepsilon^{N-1}} \left(\frac{\beta_H}{\beta}\right)^N. \quad (32)$$

Notice that the last expression (32) includes no explicit dependence on the parameters of the charged dilatonic black hole, r_+ , r_- , and the dilaton coupling a . Thus the same expression holds for a Schwarzschild or a Reissner-Nordström black hole. Our result is in accord with the one in Refs. [2] and [5] and the other reports for $N = 3$. Moreover, since the entropy at the “tree level” is also proportional to the “area” of the horizon surface, i.e.,

$$S_0 = \frac{1}{4} \tilde{A}_H^{(N-1)}, \quad (33)$$

the contribution (32) to the entropy can be treated by “renormalization” of the gravitational constant [5, 8].

Next, we turn to the subleading divergences (as $\varepsilon \rightarrow 0$). The divergences of the order of $\varepsilon^{-(N-3)}$ ($N \neq 3$) is present in both the terms of $O(\beta^{-(N+1)})$ and $O(\beta^{-(N-1)})$. For $N = 3$, these divergences are proportional to the logarithm of ε .

For $N \neq 3$, we find

$$\begin{aligned} & F_{div2} \\ = & -\frac{(N-2)A_{N-1}}{12(N-3)(2\pi)^N} \frac{\Gamma((N+1)/2)\zeta(N+1)}{\pi^{(N+1)/2}} \frac{\tilde{r}_+^{N-3}}{\varepsilon^{N-3}} \frac{\beta_H^N}{\beta^{N+1}} \\ & \times \left[(N-1)(N+3) \left(1 - \frac{r_-^{N-2}}{r_+^{N-2}}\right) + \left(\frac{(3N^2 - 2N - 3)a^2}{N-2+a^2} - 2N(N-2)\right) \frac{r_-^{N-2}}{r_+^{N-2}} \right] \\ & - \frac{A_{N-1}}{4(N-3)(2\pi)^{N-2}} \frac{\Gamma((N-1)/2)\zeta(N-1)}{\pi^{(N+1)/2}} \frac{\tilde{r}_+^{N-3}}{\varepsilon^{N-3}} \frac{\beta_H^{N-2}}{\beta^{N-1}} \\ & \times \left[\frac{(N-1)(N-2)}{6} - \frac{(N-1)(N-2)}{2} \left(1 - \frac{r_-^{N-2}}{r_+^{N-2}}\right) - m^2 \tilde{r}_+^2 \sigma^{2(b-c)}(r_+) \right], \quad (34) \end{aligned}$$

and for $N = 3$ we obtain

$$\begin{aligned} F_{div2} = & -\frac{1}{720} \frac{\beta_H^3}{\beta^4} \left[2 - \frac{3}{1+a^2} \frac{r_-}{r_+} \right] \ln \left(\frac{\Lambda^2}{\varepsilon^2} \right) \\ & - \frac{1}{72} \frac{\beta_H}{\beta^2} \left[1 - 3 \left(1 - \frac{r_-}{r_+}\right) - 3m^2 \tilde{r}_+^2 \sigma^{2(b-c)}(r_+) \right] \ln \left(\frac{\Lambda^2}{\varepsilon^2} \right), \quad (35) \end{aligned}$$

where Λ is IR cutoff.

These lead to the following contributions to the entropy:

$$\begin{aligned}
& S_{div2} \\
&= \frac{(N+1)(N-2)A_{N-1}}{12(N-3)(2\pi)^N} \frac{\Gamma((N+1)/2)\zeta(N+1)}{\pi^{(N+1)/2}} \frac{\tilde{r}_+^{N-3}}{\varepsilon^{N-3}} \left(\frac{\beta_H}{\beta}\right)^N \\
&\times \left[(N-1)(N+3) \left(1 - \frac{r_-^{N-2}}{r_+^{N-2}}\right) + \left(\frac{(3N^2 - 2N - 3)a^2}{N-2+a^2} - 2N(N-2)\right) \frac{r_-^{N-2}}{r_+^{N-2}} \right] \\
&+ \frac{(N-1)A_{N-1}}{4(N-3)(2\pi)^{N-2}} \frac{\Gamma((N-1)/2)\zeta(N-1)}{\pi^{(N+1)/2}} \frac{\tilde{r}_+^{N-3}}{\varepsilon^{N-3}} \left(\frac{\beta_H}{\beta}\right)^{N-2} \\
&\times \left[\frac{(N-1)(N-2)}{6} - \frac{(N-1)(N-2)}{2} \left(1 - \frac{r_-^{N-2}}{r_+^{N-2}}\right) - m^2 \tilde{r}_+^2 \sigma^{2(b-c)}(r_+) \right], \quad (36)
\end{aligned}$$

for $N \neq 3$, and

$$\begin{aligned}
S_{div2} &= \frac{1}{180} \left(\frac{\beta_H}{\beta}\right)^3 \left[2 - \frac{3}{1+a^2} \frac{r_-}{r_+} \right] \ln \left(\frac{\Lambda^2}{\varepsilon^2}\right) \\
&+ \frac{1}{36} \frac{\beta_H}{\beta} \left[1 - 3 \left(1 - \frac{r_-}{r_+}\right) - 3m^2 \tilde{r}_+^2 \sigma^{2(b-c)}(r_+) \right] \ln \left(\frac{\Lambda^2}{\varepsilon^2}\right), \quad (37)
\end{aligned}$$

for $N = 3$.

This result for $N = 3$ can be compared with the results of other authors. The first line of Eq. (37) agrees with the evaluation of Ghosh and Mitra [7], who also adopted the WKB method, when $a = 1$ and in the limit $r_- = r_+$ after setting $\beta = \beta_H$.

The subleading contribution to the entropy has been evaluated also by Solodukhin [8]. Their logarithmic terms for Schwarzschild and Reissner-Nordström black holes (Eqs. (3) and (20) in the second paper in [8]) coincide with the first term in (37) with $a = 0$. For a dilatonic black hole, although the expressions do not coincide exactly, the qualitative feature (its positivity in the extremal limit) is similar. The origin of the difference is not clear up till now.

The second line in the expression (37) has been found in this paper for the first time. This may not be a small contribution in comparison with the rest. Although we have only given the first and second terms in the expansion, we would get the general feature of the series of divergences. For a charged dilatonic black hole (with $a \neq 0$) in the extreme limit $r_- \rightarrow r_+$ (after setting $\beta = \beta_H$), the divergent quantum corrections as $\varepsilon \rightarrow 0$ would vanish except for a logarithmic term (for odd N), since higher divergent contribution would be written in positive powers of $\tilde{r}_+$.

To summarise, we have computed the leading and subleading divergence in the quantum correction to black hole entropy by using the WKB approximation. We have confirmed that the leading divergent piece of the entropy is proportional to the “area” of the horizon and has no explicit dependence on other parameters describing the black hole. The subleading divergence obtained in this paper coincides partially with the known result. We have shown that there is a piece

of the subleading divergence that comes from the next-to-leading term in the high-temperature expansion.

References

- [1] G. W. Gibbons and S. W. Hawking, Phys. Rev. **D15**, 2752 (1977).
- [2] G. 't Hooft, Nucl. Phys. **B256**, 727 (1985).
- [3] C. G. Callan and F. Wilczek, Phys. Lett. **B333**, 55 (1994); J. S. Dowker, Class. Quantum Grav. **11**, L55 (1994); C. Holzhey, F. Larsen and F. Wilczek, Nucl. Phys. **B424**, 443 (1994); F. Larsen and F. Wilczek, Ann. Phys. **243**, 280 (1995); R. Emparan, Phys. Rev. **D51**, 5716 (1995); M. McGuigan, Phys. Rev. **D50**, 5225 (1994).
- [4] L. Bombelli, R. Koul, J. Lee and R. Sorkin, Phys. Rev. **D34**, 373 (1986); M. Srednicki, Phys. Rev. Lett. **71**, 666 (1993).
- [5] L. Susskind and J. Uglum, Phys. Rev. **D50**, 2700 (1994).
- [6] D. Kabat and M. J. Strassler, Phys. Lett. **B329**, 46 (1994).
- [7] A. Ghosh and P. Mitra, Phys. Rev. Lett. **73**, 2521 (1994).
- [8] S. N. Solodukhin, Phys. Rev. **D51**, 609 (1995); Phys. Rev. **D51**, 618 (1995).
- [9] G. W. Gibbons and K. Maeda, Nucl. Phys. **B298**, 741 (1988); D. Garfinkle, G. T. Horowitz and A. Strominger, Phys. Rev. **D43**, 3140 (1991).
- [10] K. Shiraishi, Mod. Phys. Lett. **A7**, 3449 (1992).
- [11] J. L. F. Barbon, Phys. Rev. **D50**, 2712 (1994); Phys. Lett. **B339**, 41 (1994).
- [12] D. Birmingham, R. Kantowski and K. A. Milton, Phys. Rev. **D38**, 1809 (1988).
- [13] Eds. M. Abramowitz and I. A. Stegun, *Handbook of Mathematical Functions* (Dover, 1972).